

CORRECTION

Open Access



Correction to: Foraging conditions for breeding penguins improve with distance from colony and progression of the breeding season at the South Orkney Islands

Jessica Ann Phillips^{1*} , Annette L. Fayet¹, Tim Guilford¹, Fabrizio Manco², Victoria Warwick-Evans³ and Phil Trathan^{3*}

Correction to: *Movement Ecology* (2021) 9:22

<https://doi.org/10.1186/s40462-021-00261-x>

Following publication of the original article [1], it was noted that due to a typesetting error, Figs. 3 and 4 were swapped and paired with the wrong caption. The correct

Figs. 3 and 4 and their captions have been included in this Correction and the original article has been corrected.

The publisher apologises to the authors and readers for the inconvenience caused by the error.

The original article can be found online at <https://doi.org/10.1186/s40462-021-00261-x>

*Correspondence: jessy.a.phillips@gmail.com; pnt@bas.ac.uk

¹ Department of Zoology, Oxford University, 11a Mansfield Rd, Oxford OX1 3SZ, UK

³ British Antarctic Survey, High Cross, Madingley Rd, Cambridge CB3 0ET, UK

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 3 Depth of chinstrap penguin foraging dives during incubation and brood at four colonies in the South Orkney Islands




Fig. 4 Maximum distance from the colony of foraging trips by start date during incubation and brood at **a)** Laurie Island, **b)** Powell Island, **c)** Monroe Island, and **d)** Signy Island. The shaded areas represent slope standard error

Author details

¹Department of Zoology, Oxford University, 11a Mansfield Rd, Oxford OX1 3SZ, UK. ²Anglia Ruskin University, Cambridge Campus, East Rd, Cambridge CB1 1PT, UK. ³British Antarctic Survey, High Cross, Madingley Rd, Cambridge CB3 0ET, UK.

Published online: 04 August 2022

Reference

- Phillips JA, et al. Foraging conditions for breeding penguins improve with distance from colony and progression of the breeding season at the South Orkney Islands. *Mov Ecol.* 2021;9:22. <https://doi.org/10.1186/s40462-021-00261-x>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.